arXiv:1611.06577v1 [math.NT] 20 Nov 2016

Shifted Character Sums with Multiplicative Coefficients, II

K. Gong, C. Jia and M. A. Korolev

Abstract. Let $f(n)$ be a multiplicative function with $|f(n)| \leq 1$, q be a prime number and a be an integer with $(a, q) = 1$, χ be a non-principal Dirichlet character modulo q. Let ε be a sufficiently small positive constant, A be a large constant, $q^{\frac{1}{2}+\varepsilon} \ll N \ll q^A$. In this paper, we shall prove that

$$\sum_{n\leq N} f(n)\chi(n+a) \ll N\frac{\log\log q}{\log q}$$

and that

$$\sum_{n\leq N} f(n)\chi(n+a_1)\cdots\chi(n+a_t) \ll N\frac{\log\log q}{\log q},$$

where $t \geq 2$, $a_1, \ldots, a_t$ are distinct integers modulo q.

1. Introduction

Let q be a prime number, a be an integer with $(a, q) = 1$, χ be a non-principal Dirichlet character modulo q.

Since the 1930s, I.M. Vinogradov had begun the study on character sums over shifted primes

$$\sum_{p\leq N} \chi(p+a),$$

and obtained some deep results [9], [10] where p runs through prime numbers. His best known result is a nontrivial estimate for the range $q^{\frac{3}{4}+\varepsilon} \ll N \ll q^A$, where ε is a sufficiently small positive constant and A is a large constant. Later, A.A. Karatsuba [5] widen the range to $q^{\frac{1}{2}+\varepsilon} \ll N \ll q^A$.

For the Möbius function $\mu(n)$, one can get same results on sums

$$\sum_{n\leq N} \mu(n)\chi(n+a)$$

as that on sums over shifted primes. K. Gong and C. Jia [4] considered the general sum

$$\sum_{n\leq N} f(n)\chi(n+a), \tag{1.1}$$

where $f(n)$ is a multiplicative function with $|f(n)| \le 1$. They applied a modification (see [3]) of the method in Section 2 in [2], which is called as the finite version of Vinogradov's inequality, to give a nontrivial estimate for the sum (1.1) when q is in a suitable range. Precisely, they proved that if $f(n)$ is a multiplicative function with $|f(n)| \le 1$, q is a prime number and a is an integer with $(a,\, q) = 1$, χ is a non-principal Dirichlet character modulo q, $q^{\frac{1}{2}} \le N$, then

$$\sum_{n\le N} f(n)\chi(n+a) \ll \frac{N}{q^{\frac{1}{4}}}\log\log(6N) + q^{\frac{1}{4}}N^{\frac{1}{2}}\log(6N) + \frac{N}{\sqrt{\log\log(6N)}},$$

which is nontrivial for $q^{\frac{1}{2}+\varepsilon} \ll N \ll q^A$.

In this paper, we shall use some new ideas from [7] to give a refinement on the above upper bound.

Theorem. *Let $f(n)$ be a multiplicative function with $|f(n)| \le 1$, q be a prime number and a be an integer with $(a,\, q) = 1$, χ be a non-principal Dirichlet character modulo q. Let ε be a sufficiently small positive constant, A be a large constant, $q^{\frac{1}{2}+\varepsilon} \ll N \ll q^A$. Then we have*

$$\sum_{n\le N} f(n)\chi(n+a) \ll N\frac{\log\log q}{\log q}, \tag{1.2}$$

where the implied constant depends on ε, A.

For $t \ge 2$ and distinct integers $a_1, \ldots, a_t$ modulo q, we have

$$\sum_{n\le N} f(n)\chi(n+a_1)\cdots\chi(n+a_t) \ll N\frac{\log\log q}{\log q}, \tag{1.3}$$

where implied constant depends on ε, A and t.

Throughout this paper, we assume that q is a sufficiently large prime number, ε is a sufficiently small positive constant, A is a large constant. Let p denote prime number. Set

$$X = \log^2 q, \qquad Y = q^{\frac{\varepsilon}{4}}. \tag{1.4}$$

2. The proof of Theorem

We need one auxiliary assertion. Write

$$R = \{n :\, n \le N,\, n \text{ has a prime factor in } (X,\, Y]\},$$
$$T = \{n :\, n \le N,\, n \text{ has no prime factor in } (X,\, Y]\}.$$

Lemma 1. *We have*

$$|T| \ll N\frac{\log\log q}{\log q}.$$

Proof. Let $\Phi(x,y)$ be the number of $n \le x$ free of prime divisors $\le y$ and denote by $\Psi(x,y)$ the number of $n \le x$ free of prime divisors $> y$. Then, uniformly in $2 \le y \le x$, we have

$$\Phi(x,y) \ll \frac{x}{\log y}, \quad \Psi(x,y) \ll xe^{-\frac{u}{2}}, \quad u = \frac{\log x}{\log y}. \tag{2.1}$$

The proof of these inequalities can be found in [6] and in [8, Part III, Ch. III.5, §5.1] correspondingly.

Any $n \in T$ has the form uv: (a) all prime divisors of u do not exceed X (or $u = 1$), and all prime divisors of v are greater than Y; (b) all prime divisors of u do not exceed X (or $u = 1$) and $v = 1$. If N_a and N_b denote the number of $n \in T$ satisfying (a) and (b) respectively, then $|T| \le N_a + N_b$.

Fixing the factor u, we obtain at most $\Phi(Nu^{-1}, Y)$ possibilities to choose the factor $v = nu^{-1} \le Nu^{-1}$. If $v \ne 1$, then $v > Y$ and hence $Nu^{-1} > Y$. By (2.1), we get

$$\Phi\left(\frac{N}{u}, Y\right) \ll \frac{N}{\log Y} \cdot \frac{1}{u}.$$

The summation over u yields

$$\begin{aligned} N_a &\ll \frac{N}{\log Y}\sum_u \frac{1}{u} \ll \frac{N}{\log Y}\prod_{p\le X}\left(1+\frac{1}{p}+\frac{1}{p^2}+\cdots\right) \\ &\ll \frac{N}{\log Y}\prod_{p\le X}\left(1-\frac{1}{p}\right)^{-1} \ll N\frac{\log X}{\log Y} \ll N\frac{\log\log q}{\log q}. \end{aligned}$$

Using (2.1) again, we obtain

$$N_b \le \Psi(N,X) \ll N\exp\left(-\frac{\log N}{2\log X}\right) \ll N\frac{\log X}{\log N} \ll N\frac{\log\log q}{\log q}.$$

Hence, Lemma 1 is proved.

Now we begin the proof of Theorem. The discussion in Lemma 1 of [4] yields

$$\sum_{n\le N} f(n)\chi(n+a) = \sum_{\substack{n\le N\\(n,q)=1}} f(n)\chi(n+a) + O\Big(\frac{N}{q}+1\Big).$$

By Lemma 1,

$$\sum_{\substack{n\le N\\(n,\,q)=1}} f(n)\chi(n+a) = \sum_{\substack{n\le N\\ n\in R\\(n,\,q)=1}} f(n)\chi(n+a) + O\Big(N\frac{\log\log q}{\log q}\Big).$$

Therefore we have

$$\sum_{n\le N} f(n)\chi(n+a) = \sum_{\substack{n\le N\\ n\in R\\(n,\,q)=1}} f(n)\chi(n+a) + O\Big(N\frac{\log\log q}{\log q}\Big). \tag{2.2}$$

The contribution of terms in which n is divisible by p^2 for some p under the condition $X < p \le Y$ is

$$\ll \sum_{X<p\le Y} \frac{N}{p^2} \ll \frac{N}{\log^2 q}.$$

Denote by B_r the set of $n \le N$ not dividing by p^2 for some $p \in (X, Y]$ and having exactly r distinct prime factors in $(X, Y]$. Then

$$\sum_{\substack{n\le N\\ n\in R\\(n,\,q)=1}} f(n)\chi(n+a) = \sum_{1\le r\ll \log N} \sum_{\substack{n\le N\\ n\in B_r\\(n,\,q)=1}} f(n)\chi(n+a) \; + O\Big(\frac{N}{\log^2 q}\Big). \tag{2.3}$$

Among the products pm ($pm \le N$, $(p,\, m) = 1$, $X < p \le Y$, $m \in B_{r-1}$), the number $n(\in B_r)$ appears r times. Thus

$$\sum_{\substack{n\le N\\ n\in B_r\\(n,\,q)=1}} f(n)\chi(n+a) = \frac{1}{r}\sum_{\substack{X<p\le Y\\(p,\,q)=1}} f(p) \sum_{\substack{m\in B_{r-1}\\ mp\le N\\(m,\,p)=1\\(m,\,q)=1}} f(m)\chi(pm+a).$$

If we get rid of the condition $(m,\, p) = 1$, then the corresponding error is

$$\ll \frac{1}{r}\sum_{X<p\le Y}\sum_{\substack{m\le \frac{N}{p}\\ p\mid m}} 1 \ll \frac{1}{r}\sum_{X<p\le Y}\frac{N}{p^2} \ll \frac{1}{r}\sum_{X<p}\frac{N}{p^2} \ll \frac{N}{rX}.$$

Therefore we have

$$\sum_{\substack{n\le N\\ n\in B_r\\(n,\,q)=1}} f(n)\chi(n+a) = \frac{1}{r}\sum_{\substack{X<p\le Y\\(p,\,q)=1}} f(p) \sum_{\substack{m\in B_{r-1}\\ mp\le N\\(m,\,q)=1}} f(m)\chi(pm+a) + O\Big(\frac{N}{rX}\Big).$$

We divide $\{X < p \le Y\}$ into $O(\log Y)$ intervals of the form $\{Q < p \le Q_1\}$, where $X \le Q \le Q_1 \le Y$, and write

$$\begin{aligned} S_r(Q,\, Q_1) &= \sum_{\substack{Q<p\le Q_1\\(p,\,q)=1}} f(p) \sum_{\substack{m\in B_{r-1}\\ mp\le N\\(m,\,q)=1}} f(m)\chi(pm+a) \\ &= \sum_{\substack{m\in B_{r-1}\\ m\le \frac{N}{Q}\\(m,\,q)=1}} f(m) \sum_{\substack{Q<p\le Q_1\\ p\le \frac{N}{m}\\(p,\,q)=1}} f(p)\chi(pm+a). \end{aligned}$$

We have

$$|S_r(Q,\, Q_1)| \le \sum_{m\le \frac{N}{Q}} \Big| \sum_{\substack{Q<p\le Q_1\\ p\le \frac{N}{m}\\(p,\,q)=1}} f(p)\chi(pm+a)\Big|.$$

We shall estimate the sum

$$W_1 = \sum_{m\le \frac{N}{Q}} \Big| \sum_{\substack{Q<p\le Q_1\\ p\le \frac{N}{m}\\(p,\,q)=1}} f(p)\chi(pm+a)\Big|. \tag{2.4}$$

In the following, we need

Lemma 2. *Let q be a prime number, χ be a non-principal Dirichlet character modulo q, $(a,\, q) = 1$. Then for two primes p_1, p_2 with $(p_1p_2,\, q) = 1$, $p_1 \not\equiv p_2 \,(\mathrm{mod}\, q)$, we have*

$$\sum_{U<m\le V} \chi\Big(\frac{p_1m+a}{p_2m+a}\Big) \ll \frac{V-U}{\sqrt{q}} + \sqrt{q}\log q.$$

Here we write $\frac{1}{n}$ as the multiplicative inverse of n such that $\frac{1}{n}\cdot n \equiv 1 \,(\mathrm{mod}\, q)$ and appoint $\frac{1}{0} = 0$.

This is Lemma 4 in [4] which is produced by Lemma 18 in [1].

By Cauchy's inequality, we have

$$|W_1|^2 \le \frac{N}{Q} \sum_{m\le \frac{N}{Q}} \Big| \sum_{\substack{Q<p\le Q_1\\ p\le \frac{N}{m}\\(p,\,q)=1}} f(p)\chi(pm+a)\Big|^2.$$

Then

$$\begin{aligned}
W_2 &= \sum_{m\le \frac{N}{Q}} \Big| \sum_{\substack{Q<p\le Q_1\\ p\le \frac{N}{m}\\ (p,\,q)=1}} f(p)\chi(pm+a)\Big|^2 \\
&= \sum_{m\le \frac{N}{Q}} \sum_{\substack{Q<p_1\le Q_1\\ p_1\le \frac{N}{m}\\ (p_1,\,q)=1}} \sum_{\substack{Q<p_2\le Q_1\\ p_2\le \frac{N}{m}\\ (p_2,\,q)=1}} f(p_1)\overline{f(p_2)}\chi\Big(\frac{p_1m+a}{p_2m+a}\Big) \\
&= \sum_{\substack{Q<p_1\le Q_1\\ (p_1,\,q)=1}} \sum_{\substack{Q<p_2\le Q_1\\ (p_2,\,q)=1}} f(p_1)\overline{f(p_2)} \sum_{\substack{m\le \frac{N}{Q}\\ m\le \frac{N}{\max(p_1,\,p_2)}}} \chi\Big(\frac{p_1m+a}{p_2m+a}\Big) \\
&\ll \sum_{\substack{Q<p_1\le Q_1\\ (p_1,\,q)=1}} \sum_{\substack{p_1\le p_2\le Q_1\\ (p_2,\,q)=1}} \Big| \sum_{m\le \frac{N}{p_2}} \chi\Big(\frac{p_1m+a}{p_2m+a}\Big)\Big| \\
&\ll \sum_{Q<p_1\le Q_1} \frac{N}{Q} + \sum_{\substack{Q<p_1\le Q_1\\ (p_1,\,q)=1}} \sum_{\substack{p_1<p_2\le Q_1\\ (p_2,\,q)=1}} \Big| \sum_{m\le \frac{N}{p_2}} \chi\Big(\frac{p_1m+a}{p_2m+a}\Big)\Big|.
\end{aligned}$$

We note that if $p_1,\, p_2 \le Y = q^{\frac{\varepsilon}{4}}$, then $p_1 \ne p_2$ is equivalent to $p_1 \not\equiv p_2 \,(\mathrm{mod}\, q)$. By Lemma 2,

$$\begin{aligned}
W_2 &\ll N + \sum_{Q<p_1\le Q_1} \sum_{Q<p_2\le Q_1} \Big(\frac{N}{\sqrt{q}Q} + \sqrt{q}\log q\Big) \\
&\ll N + \frac{NQ}{\sqrt{q}} + Q^2\sqrt{q}\log q \\
&\ll N.
\end{aligned}$$

Hence,

$$|S_r(Q,\, Q_1)| \le W_1 \ll \frac{N}{\sqrt{Q}}.$$

Taking

$$Q = 2^k X, \qquad k = 0,\, 1,\, \ldots,\, k_0, \qquad 2^{k_0-1}X \le Y < 2^{k_0}X,$$

we get

$$\sum_{\substack{X<p\le Y\\ (p,\,q)=1}} f(p) \sum_{\substack{m\in B_{r-1}\\ mp\le N\\ (m,\,q)=1}} f(m)\chi(pm+a) \ll N\sum_{k=0}^{k_0} \frac{1}{2^{\frac{k}{2}}\sqrt{X}} \ll \frac{N}{\log q}.$$

It follows that

$$\sum_{\substack{n\le N\\ n\in B_r\\ (n,\,q)=1}} f(n)\chi(n+a) \ll \frac{N}{r\log q}.$$

Then

$$\sum_{\substack{n\le N\\ n\in R\\ (n,\,q)=1}} f(n)\chi(n+a) \ll \sum_{1\le r\ll \log N} \frac{N}{r\log q} \ll N\frac{\log\log q}{\log q}.$$

Thus

$$\sum_{n\le N} f(n)\chi(n+a) \ll N\frac{\log\log q}{\log q}.$$

We have proved (1.2) for sufficiently large prime number q. For bounded q, this is a trivial result. The proof of (1.3) follows the same lines, but we need to replace Lemma 2 in the above arguments by the following

Lemma 3. *Let q be a prime number, χ be a non-principal Dirichlet character modulo q. Assume that $t \ge 2$, $(b_1,\, q) = \cdots = (b_t,\, q) = 1$, $b_1, \ldots, b_t$ are distinct integers modulo q and that primes p_1, p_2 satisfy $(p_1p_2,\, q) = 1$, $p_1 \not\equiv p_2 \pmod q$. Then if $p_2 \not\equiv \overline{b_i}b_jp_1 \pmod q$ $(1 \le i,\, j \le t)$, we have*

$$\sum_{U<m\le V} \chi\Big(\frac{(p_1m+b_1)\cdots(p_1m+b_t)}{(p_2m+b_1)\cdots(p_2m+b_t)}\Big) \ll \frac{V-U}{\sqrt{q}} + \sqrt{q}\log q.$$

This is Lemma 6 in [4] which is produced by Lemma 17 in [1]. Lemma 17 in [1] is due to A. Weil [11, Appendix V, Example 12], which states that

$$\left|\sum_{x=1}^{q} \chi_1(x+c_1)\cdots\chi_r(x+c_r)e^{2\pi i\,\frac{f(x)}{q}}\right| \le\ (r+d)\sqrt{q}.$$

Here $\chi_1, \ldots, \chi_r$ are any Dirichlet characters modulo q such that at least one of them is non-principal, $f \in \mathbb{F}_q[x]$ is any polynomial of degree $d \ge 0$ and $c_1, \ldots, c_r$ are distinct integers modulo q.

We should discuss in two cases respectively. In the first case, there is no a_i in (1.3) is equivalent to 0 modulo q. In the second case, one of a_i's is equivalent to 0 modulo q, which is reduced to the case of $t-1$ by writing $f(n)\chi(n) = f_1(n)$. Then the proof of (1.3) follows.

So far the proof of Theorem is complete.

Acknowledgements

The first author is supported by the National Natural Science Foundation of China (Grant No. 11671119). The second author is supported by the National Key Basic Research Program of China (Project No. 2013CB834202) and the National Natural Science Foundation of China (Grant No. 11371344 and Grant No. 11321101). The work of the last author is supported by Russian Science Foundation (Grant No. 14-11-00433) and performed in Steklov Mathematical Institute of Russian Academy of Sciences.

References

[1] J. Bourgain, M.Z. Garaev, S.V. Konyagin and I.E. Shparlinski, *On the hidden shifted power problem*, SIAM J. Comput. **41** (2012), 1524–1557.

[2] J. Bourgain, P. Sarnak and T. Ziegler, *Disjointness of Mobius from horocycle flows*, From Fourier Analysis and Number Theory to Radon Transforms and Geometry, 67–83, Dev. Math. **28**, Springer, New York, 2013.

[3] K. Gong and C. Jia, *Kloosterman sums with multiplicative coefficients*. Sci. China Math. **59** (2016), 653–660.

[4] K. Gong and C. Jia, *Shifted character sums with multiplicative coefficients*, J. Number Theory **153** (2015), 364–371.

[5] A.A. Karatsuba, *Sums of characters over prime numbers*, Izv. Akad. Nauk SSSR Ser. Mat. **34** (1970), 299–321 (Russian); English translation in Math. USSR Izv. **4** (1970), 303–326.

[6] M.A. Korolev, *On Karatsuba's method of estimating Kloosterman sums*, Mat. Sb. **207** (2016), 117–134.

[7] M.A. Korolev, *On Kloosterman sums with multiplicative coefficients*, `arXiv:1610.09171 [math.NT]`.

[8] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, Cambridge Univ. Press, Cambridge, 1995.

[9] I.M. Vinogradov, *On the distribution of quadratic rests and non-rests of the form $p + k$ to a prime modulus*, Mat. Sb. **3** (1938), 311–319. (Russian)

[10] I.M. Vinogradov, *Improvement of an estimate for the sum of the values $\chi(p+k)$*, Izv. Akad. Nauk SSSR Ser. Mat. **17** (1953), 285–290. (Russian)

[11] A. Weil, *Basic Number Theory*, Springer-Verlag, New York, Heidelberg, Berlin, 1974.

Ke Gong

Department of Mathematics, Henan University, Kaifeng, Henan 475004, P. R. China

E-mail: kg@henu.edu.cn

Chaohua Jia

Institute of Mathematics, Academia Sinica, Beijing 100190, P. R. China

Hua Loo-Keng Key Laboratory of Mathematics, Chinese Academy of Sciences, Beijing 100190, P. R. China

E-mail: jiach@math.ac.cn

Maxim Aleksandrovich Korolev

Steklov Mathematical Institute of Russian Academy of Sciences

119991, Russia, Moscow, Gubkina Str., 8

E-mail: korolevma@mi.ras.ru